



# OPAC-2010 Workshop

September 8, Graz, Austria



## Consistency of Radio Occultation Climatologies from Different Satellites

U. Foelsche, B. Scherllin-Pirscher\*, A. Leuprecht,  
and G. Kirchengast,

WegCenter and IGAM/IP, University of Graz, Austria

\* now at UCAR, Boulder, CO, USA

funded by





# Acknowledgements



We would like to thank the following centers for fruitful collaboration and for the provision of excess phase and orbit data:

**UCAR** (University Corporation for Atmospheric Research) in Boulder, CO, USA

**GFZ** (German Research Centre for Geosciences) Potsdam, Germany

**EUMETSAT** (European Organisation for the Exploitation of Meteorological Satellites) in Darmstadt, Germany

**ECMWF** (European Centre for Medium-Range Weather Forecasts), Reading, UK, is acknowledged for the continuous provision of analysis and forecast data



# Motivation



Long-term climate monitoring based on radio occultation data will require the combination of data from different satellites.

In the best case we would hope for:

**Consistent results** for data from **different satellites**

Consistent results for retrievals from **different centers**  
(see talk by Andrea Steiner)



# WegCenter RO Profiles



Profile retrieval, termed “**Occultation Processing System – OPS**” (current version: 5.4) starts with **excess phase** and **orbit data** from different processing centers (UCAR, GFZ, EUMETSAT, ..).

**Consistent retrieval for all satellites.**

High altitude initialization by:

**Statistical Optimization** using **ECMWF forecasts**.

OPsv5.4 uses (still) geometric optics and dry air retrieval, focusing on the UTLS.

“**Occultation event data**” (OED), **Level 2: Single Profiles**, format, including **Retrieval to A Priori Error Ratio (RAER)** and **Tropopause Parameters**.




# WegCenter Climatologies



## Binning and Averaging – “Climate Processing System” (CLIPS)

- Radio Occultation profiles in a “bin” are collected and averaged
- “**Fundamental**” bins with **5° latitudinal** and **60° longitudinal** width are combined to build zonal mean **monthly** climatologies.
- **Basic latitudinal resolution** is **10°**, each of the 18 latitude bands (pole to pole) contains **two 5° bands**, and the mean profiles for these two bands are averaged, weighted with the surface area.
- **Seasonal** climatologies are obtained by averaging over three months.
- **Annual** climatologies by averaging over a **calendar year**.
- 200 m vertical spacing was chosen for the altitude gridding.

**Climatology Data (CLD), Level 3:** Bending Angle, Refractivity, Pressure, Geopotential Height, Dry Temperature.



Different sampling strategies, depending on the latitudinal distribution


**Straight line** (equal numbers **per latitude**): **cosine-weighting** (CHAMP)

**Curved line** (equal numbers **per area**): **No weighting** (COSMIC-II ?)


Currently: cosine-weighting for all satellites

# Data Quality


CHAMP:	Mean: $-0.16 \mu\text{rad}$
GRACE-A:	Mean: $-0.12 \mu\text{rad}$
F3C-FM1:	Mean: $-0.14 \mu\text{rad}$
F3C-FM2:	Mean: $-0.14 \mu\text{rad}$
F3C-FM3:	Mean: $-0.14 \mu\text{rad}$
F3C-FM4:	Mean: $-0.14 \mu\text{rad}$
F3C-FM5:	Mean: $-0.14 \mu\text{rad}$
F3C-FM6:	Mean: $-0.14 \mu\text{rad}$
MetOp:	Mean: $-0.09 \mu\text{rad}$




CHAMP:	Mean: $4.74 \mu\text{rad}$
GRACE-A:	Mean: $2.70 \mu\text{rad}$
F3C-FM1:	Mean: $2.65 \mu\text{rad}$
F3C-FM2:	Mean: $2.46 \mu\text{rad}$
F3C-FM3:	Mean: $2.62 \mu\text{rad}$
F3C-FM4:	Mean: $2.54 \mu\text{rad}$
F3C-FM5:	Mean: $2.71 \mu\text{rad}$
F3C-FM6:	Mean: $2.49 \mu\text{rad}$
MetOp:	Mean: $0.98 \mu\text{rad}$




**Bending Angle Statistics between 65 km and 80 km**  
 “Bias” means – difference to **MSIS** climatology.  
 Annual and **semi-annual cycle**.









# Single-COSMIC Climatology



DJF 2007/08 dry temperature **variability**. Mean state is much harder to capture, if the variability is high, e.g., **tropopause** region, **winter Extratropics** vs. Low Lat

# Seasonal F3C-FM4 Climatology



# Sampling Error


Even with perfect observations, climatologies from two satellites would be slightly different, because measurements are taken at different times and locations

**Estimation** of the sampling error based on **ECMWF** 3D fields on four time layers per day (as proxy for the “true” temporal and spatial evolution):


“True” **profiles** at the RO locations  
minus  
“True” **mean field**

# Consistency of RO Data

Tdry: Year 2007: F3C/FM-1 - F3C/FM-4



Tdry: Year 2007: (F3C/FM-1-Sampl Err) - (F3C/FM-4-Sampl Err)




Systematic Differences between 2007 **annual mean**, zonal mean Tdry climatologies from different **Formosat-3/ COSMIC** satellites (FM-1 and FM-4).

After **subtraction of the estimated sampling errors** from both climatologies, differences are  $< 0.1$  K almost everywhere (despite errors in sampling error estimation for both satellites)

# Consistency – JJA 2007

COSMIC-FM2 minus COSMIC-FM4


Tdry: JJA 2007: (F3C/FM-2-Sampl Err) - (F3C/FM-4-Sampl Err)




# Consistency of RO Data

Oct 2007: Deviations of Dry Temperature (Sampling Error Subtracted)


CHAMP-Satellite Mean




GRACE-A-Satellite Mean



F3C/FM-1-Satellite Mean






F3C/FM-2-Satellite Mean



Monthly mean dry temperature (Oct 2007) wrt satellite mean

# Consistency of RO Data




**GRAS** excess phase data have only been available in offline mode. There seems to be something wrong in our Tdry retrievals


Monthly mean dry temperature (Oct 2007) wrt satellite mean

# Refractivity Consistency

CHAMP:	Mean: 0.01 %
GRACE-A:	Mean: -0.00 %
F3C/FM-1:	Mean: -0.00 %
F3C/FM-2:	Mean: 0.01 %
F3C/FM-3:	Mean: -0.03 %
F3C/FM-4:	Mean: 0.01 %
F3C/FM-5:	Mean: 0.01 %
F3C/FM-6:	Mean: 0.01 %



CHAMP:	Mean: -0.01 %
GRACE-A:	Mean: -0.00 %
F3C/FM-1:	Mean: 0.00 %
F3C/FM-2:	Mean: 0.00 %
F3C/FM-3:	Mean: 0.00 %
F3C/FM-4:	Mean: -0.00 %
F3C/FM-5:	Mean: 0.00 %
F3C/FM-6:	Mean: 0.00 %




Monthly **refractivity** differences relative to the **satellite mean**


Further reduced when the estimated **sampling errors** are **subtracted**

**CHAMP** “excursion” (MAM 2007) corresponds to high BA noise

CHAMP:	Mean: -0.00 K
GRACE-A:	Mean: -0.01 K
F3C/FM-1:	Mean: 0.00 K
F3C/FM-2:	Mean: 0.01 K
F3C/FM-3:	Mean: 0.03 K
F3C/FM-4:	Mean: -0.02 K
F3C/FM-5:	Mean: 0.00 K
F3C/FM-6:	Mean: -0.01 K



CHAMP:	Mean: -0.07 K
GRACE-A:	Mean: -0.00 K
F3C/FM-1:	Mean: 0.01 K
F3C/FM-2:	Mean: 0.01 K
F3C/FM-3:	Mean: 0.00 K
F3C/FM-4:	Mean: 0.01 K
F3C/FM-5:	Mean: 0.00 K
F3C/FM-6:	Mean: 0.01 K



Monthly **dry temperature** differences relative to the **satellite mean**  
 Further reduced when the estimated **sampling errors** are **subtracted**  
**CHAMP** offset mainly between 25 km and 30 km



# Summary



- RO data from different satellites are encouragingly **consistent**
- Even more, when the **sampling errors** of both climatologies are subtracted
- Different measurement **noise** leads to different downward propagation of potential biases in the background, used for statistical optimization
- For Tdry more pronounced than for refractivity

[www.globclim.org](http://www.globclim.org)



**Thank you!**