



wege entstehen, indem wir sie gehen
paths emerge in that we walk them

Wegener Center
www.wegcenter.at



Atmospheric Remote Sensing and Climate System Research Group

ARSCISys

Structural Uncertainty of the CHAMP Climate Record from Different Data Centers

A.K. Steiner ¹, S-P. Ho ², D. Hunt ², G. Kirchengast ¹, A. J. Mannucci ³,
B. Scherllin-Pirscher ^{2,1}, H. Gleisner ⁴, A. von Engel ⁵, T. Schmidt ⁶,
C. Ao ³, S. S. Leroy ⁷, U. Foelsche ¹, Y-H. Kuo ², E. R. Kursinski ⁸,
K. B. Lauritsen ⁴, S. Syndergaard ⁴, C. Marquardt ⁵, C. Rocken ²,
W. Schreiner ², S. Sokolovskiy ², J. Wickert ⁶

¹ Wegener Center for Climate and Global Change (WEGC) & IGAM/IP, Univ. Graz, Austria

² COSMIC Project Office, UCAR, Boulder, CO, USA

³ Jet Propulsion Laboratory (JPL), California Institute of Techn., Pasadena, CA, USA

⁴ Danish Meteorological Institute (DMI), Copenhagen, Denmark;

⁵ Meteorological Division, EUMETSAT, Darmstadt, Germany;

⁶ German Research Centre for Geosciences (GFZ), Potsdam, Germany

⁷ School of Engineering and Applied Sciences, Harvard Univ., Cambridge, MA, USA

⁸ Institute of Atmospheric Physics, Univ. of Arizona, Tucson, AZ, USA




- Background
- Data and study setup
- Centers and their processing
- Results
- Conclusions and outlook

- Aim: Quantifying the structural uncertainty of RO data provided by main RO processing centers

- Background:
 - 1st round study by Ho et al. (GRL, 2009)
 - Intercomparison of refractivity climatologies
 - CHAMP 2002-2006
 - 4 centers JPL, GFZ, UCAR, WEGC
 - Structural uncertainties amongst climatologies of different RO processing centers are < 0.03 %/5years for refractivity trends in large scale means

- 2nd round intercomparison studies
- Profile to profile comparison (PPC) provided by RO processing centers (study by S-P. Ho)
- Intercomparison of zonal monthly mean climatologies (MMCs)
- **CHAMP** record September 2001 to September 2008



Courtesy: J. Wickert

- Resolution/range: 5-degree zonal means, monthly means
- **Latitude regions:** Tropics 20°N–20°S
Northern/Southern mid-latitudes: 20°N/S–50°N/S
Northern/Southern high-latitudes: 50°N/S–90°N/S
- **Altitude range:** 8–30 km (200 m)
- **Altitude layers:** 8–12 km (Upper Troposphere UT)
12–16 km (Tropopause TP)
16–25 km (Lower Stratosphere LS)
25–30 km
- Focus region: Tropical UTLS
- **Parameters^a:** Bending angle $\alpha(z_a)$
Refractivity $N(z)$
Dry pressure $p_d(z)$,
Dry geopotential height $Z_d(z_p)$
Dry temperature $T_d(z)$

^a msl altitude z , impact altitude z_a ($z_a = \text{impact parameter} - \text{radius of curvature} - \text{geoid undulation}$),
pressure altitude $z_p(p)[\text{m}] = (7000 \text{ m}) \times \ln(1013.25 \text{ hPa}/p[\text{hPa}])$

Processing Center ^b	Data version	Processing Steps ^a
DMI Copenhagen, DK	OCC 18.9.353	UCAR phase & orbit data, Geometrics optics (GO), CanonTransf. (CT)<25km Optimization of α with MSIS-E90,
EUM Darmstadt, D	YAROS0.1(Beta)	UCAR phase & orbit data, GO, Optimization of N with MSISE-90
JPL Pasadena, CA, USA	v2fo_10Kp1N	Excess phase Double Differencing, CT, exponential function fit of α at 40-50km, extrapol.
UCAR Boulder, CO, USA	2009.2650	Excess phase Single Differencing, Full Spectrum Inversion in troposphere, above GO Optimization with NCAR climatology
WEGC Graz, A	OPsv5.4	UCAR phase & orbit data, GO, Statistical optimization > 30 km with ECMWF forecasts & MSIS above


^a All centers: Ionospheric correction of bending angles and dry air retrieval but different smoothing routines and quality control

^b GFZ not yet included


- **MMCs** based on provided profiles (computed by D. Hunt, UCAR)
- **Sampling error** estimation based on **ERA-Interim** for N , p_d , Z_d , T_d for 5-deg latitude bins
- **Subtraction of sampling error**: “de-sampled” 5-deg MMCs
- MMCs and de-sampled MMCs were used
- Mean difference of each center to the intercenter mean
- Seasonal means: for each center and intercenter seasonal mean
- **De-seasonalized anomaly time series**
- De-seasonalized **anomaly difference time series and trends**:
absolute anomaly differences for Z and T (anomaly – intercenter mean)
fractional anomaly differences for α , N , p :
(anomalies - intercenter seas. mean)/intercenter seas, mean*100
- 5 centers, 5 parameters, 5 latitude bins and 4 altitude layers

Number of Occultation Events

Mean no of events per 5-deg bin




No of events 5 large zonal bins over time




Results: Center's mean difference to intercentermean

DMI




- Bending Angle
- within +/- 0.1 %
- high latitudes:
DMI < 0.2 %
JPL -0.3 %






- Refractivity
- within +/- 0.1 %
- high lats > 25 km:
DMI 0.2 %
UCAR -0.3 %


EUM






UCAR



JPL




WEGC




Results: Center's mean difference to intercentermean


DMI




JPL



UCAR



WEGC



Dry Pressure (left)

- DMI: + 0.1% < 20 km (25 km tropics)
- JPL: - 0.1% < 20 km (-0.2% @ 30 km)
- UCAR: within +/- 0.1%
- WEGC: within +/- 0.1% (< 25 km)

Dry Geopotential Height (middle)

- Same pattern as pressure
- UCAR: checks troposphere


Dry Temperature (right)

- DMI: < 0.2 K tropics, 1 K @ HL > 25 km
- JPL: within -0.2 K < 20 km up to 0.5 K > 25 km
- UCAR: +/- 0.1 K except HL > 25 km
- WEGC: < 0.2 < 25 km








Results: De-seasonalized Anomaly Time Series



Wegener Center
www.wegcenter.at



Sampling error estimation




ERA-Interim, 5-deg zonal bins

ECMWF-analyses, 10-deg zonal bins


Results: Refractivity - Fract. Anomaly Differences



Wegener Center
www.wegcenter.at




Results: Refractivity De-sampled Frac.Anom. Differences




Results: Bending Angle Fract. Anomaly Difference Trends


DMI




EUM




JPL



UCAR





WEGC





- Tropics: ~ +/- 0.02 %/7yrs
- Mid-lats: ~ 0.02 %/7yrs
- TP: ~ 0.04 %/7yrs
- Hi Lats < +/- 0.05 %/7yrs up to +/0.15 %/7yrs (DMI)

Results: Refractivity Fractional Anomaly Difference Trends



DMI





EUM





JPL



UCAR



WEGC




- right panel:
saml. error subtracted
- improvement after subtraction
- Tropics: $\sim \pm 0.02$ %/7yrs
means eg.,
for LS anomaly trend =
 -0.44 ± 0.02 %/yrs
- < 25 km: < ± 0.05 %/7yrs
except DMI at NHL
- > 25 km: < ± 0.1 %/7yrs
- Basically consistent with Ho
et al. (2009)

Results: Dry Pressure/Geop. Height Anomaly Diff. Trends

Dry Pressure



Dry Geopot. Height





- Left panels: Dry pressure
- Right panels: Dry geopotential height
- Tropics & Mid-lats UTLS & High lats UT:
 - p: < 0.05 %/7yrs
 - Z: < +/- 5 m/7yrs (except UCAR for Z)
- Tropics & Mid-lats 25-30 km
 - p: < 0.1 %/7yrs
 - Z: < +/- 10 m/7yrs
- High lats 25-30 km
 - p: < 0.2 %/7yrs
 - Z: < +/- 15 m/7yrs
- E.g. LS Anomalytrend in the tropics: -0.23+/-0.03 %/yr

Results: Temperature Anomaly Difference Trends



DMI





JPL



UCAR



WEGC



- right panels: sampl. error subtracted
- Tropics UT: $\sim \pm 0.02$ K/7yrs
LS: $< \pm 0.08$ K/7yrs
- UT global & Mid-lats LS:
 $< \pm 0.1$ K/7yrs
- 25-30 km & High lats LS:
 $< \pm 0.25$ K/7yrs
(DMI -0.35 K/7yrs)
- High lats 25-30 km:
 ~ 0.5 K/7yrs
- Best performance:
tropics to mid-latitudes UTLS
- Anomaly trends:
tropical LS: 0.25 ± 0.08 K/7yrs
tropical UT: -0.54 ± 0.02 K/7yrs

CONCLUSIONS

- Structural uncertainty of RO data from 5 processing centers
- Low structural uncertainty for α and N at 8-30 km
- Consistency with Ho et al. (2009)
- Low structural uncertainty for p, Z, T in UTLS in tropics & mid-latitudes
- Higher structural uncertainty > 25 km and at high latitudes
- Reflecting different initialization methods

OUTLOOK

- Check WEGC MAM 2007
- Include GFZ data
- Estimate sampling error for 10-deg zonal bins & its uncertainty
- Estimate level 3 uncertainty – different climatology computation methods