



OPAC-3 Workshop September 2007, Graz, Austria



Climate Monitoring based on Radio Occultation Data: From CHAMP to FORMOSAT-3/COSMIC

Ulrich Foelsche^{1,2}, Barbara Pirscher¹, Michael Borsche¹,
Gottfried Kirchengast¹, and J. Wickert³

1 WegCenter and IGAM, University of Graz, Austria

2 COSMIC Program Office, UCAR, CO, USA


3 GFZ Potsdam, Germany




foelsche@ucar.edu
ulrich.foelsche@uni-graz.at



Radio Occultation with CHAMP



CHAMP: First Opportunity to create RO Climatologies



Torsten Schmidt (tschmidt@gfz-potsdam.de)
Thu Mar 2 19:34:40 2006

Torsten Schmidt (tschmidt@gfz-potsdam.de)
Thu Mar 2 19:34:40 2006

Torsten Schmidt (tschmidt@gfz-potsdam.de)
Tue Jan 30 13:51:34 2007



Observing Climate




Zonal Mean, Seasonal Climatologies: **Dry Temperature**

Foelsche et al. 2007, *Climate Dynamics*, revised

Example Season – Winter 2003/04

DJF 2003/04: CHAMP Event Distribution



Typical Winter Season
Dec-Jan-Feb (DJF) 2003/2004
Zonal mean fields
Binning and Averaging
18 latitude bands
10°latitudinal with

WegCenter Retrieval, using phase and orbit data from GFZ



WegCenter Retrieval (1)



Early outlier rejection: “ 3σ ” outlier rejection on 50 Hz sampling rate L1 and L2 phase delay data, based on a one-second moving window over the profile.

Phase delay smoothing: Smoothing of 50 Hz phase delay profiles using regularization (third order norm, regularization parameter = 10^5 , following *Syndergaard, 1999*).

Bending angle retrieval: **Geometric optics** retrieval (e.g., *Kursinski et al., 1997*)

Ionospheric correction: Linear combination of L1 and L2 **bending angles** (*Vorob'ev and Krasil'nikova, 1994*). Correction is applied to low-pass filtered bending angles (1 km moving average), L1 high-pass contribution is added after correction (*Hocke et al., 2003*). L2 bending angles <15 km derived via L1-L2 extrapolation.

Statistical optimization of bending angles: Between 30 and 120 km. Vertically correlated background (corr. length = 6 km) and observation (corr. length = 1 km) errors. Observation error estimated from observed variance of observed profile >65 km. Background error: 15%. Background information: collocated profile derived from **ECMWF** operational analysis (T42L60; resp. T42L91 as of 01/02/2006). Above ~60 km: MSISE-90 (*Hedin, 1991*).



WegCenter Retrieval (2)



Abel transform: Numerical integration over bending angle (Simpson's trapezoidal rule) from each height (impact parameter) to 120 km.

Hydrostatic integral initialization: No initialization below 120 km.
At 120 km: pressure = pressure (MSISE-90).

External quality control (for outlier profiles): Refractivity 5 km–35 km: $\Delta N/N < 10\%$; Temperature 8 km–25 km: $\Delta T < 20$ K. Reference: collocated ECMWF operational analysis profiles (T42L60 resp. T42L91 as of 01/02/2006).

Reference frame, vertical coordinate: Earth figure: WGS-84 ellipsoid; Vertical coordinate: mean-sea-level (MSL) altitude; conversion of ellipsoidal height to **MSL altitude** (at mean tangent point location) via EGM-96 geoid smoothed to $2^\circ \times 2^\circ$ resolution.

Binning and Averaging:

$$\overline{T_{\text{dry}}(z)} = \frac{1}{\sum_{i=1}^{N_{\text{prof}}(z)} \cos(\varphi_i)} \sum_{i=1}^{N_{\text{prof}}(z)} T_{\text{dry}_i}(z, \varphi_i) \cos(\varphi_i)$$

- Weighting by the cosine of the latitude.
- “Fundamental” zonal bins with **5°** latitudinal width to build zonal mean **monthly** climatologies.
- Basic latitudinal resolution is **10°**, each of the 18 latitude bands (pole to pole) contains two fundamental bins, and the mean profiles for these two bins are averaged, weighted with the surface area.
- **Seasonal** climatologies are obtained by averaging over three months.
- 200 m vertical spacing was chosen for the altitude gridding.



Climatologies



Sampling Error:

Undersampling of the true spatial
and temporal temperature
evolution



Estimation of Sampling Error







Based on ECMWF fields – 4 time layers/day
ECMWF is taken as a proxy for the “true”
temporal and spatial evolution

Sampling Error:

“True” **profiles** at the RO locations
minus
“True” **mean field**


Dry Temperature – DJF 2003/04




Systematic Difference – Tropical Tropopause

Systematic Difference – Summer


2002




2003



2004




2005



Wave-like bias structure – Antarctica

Sampling Problem – JJA2006




Summer 2006

Lack of CHAMP data
~5 weeks

“Gap” can be filled
With **GRACE** data

Sampling Error – JJA2006



Summer 2006

Lack of CHAMP data

~5 weeks

Sampling error

amazingly small


(for zonal means)

But southern (winter)

polar vortex

clearly **too warm**

Sampling Error – JJA2006



Summer 2006

CHAMP + GRACE data



Systematic Difference – JJA2006

February 2006: **Major change at ECMWF**

Horizontal Resolution: T511 → **T799**

Vertical Resolution: L60 → **L91**

Top model level: 0.1 hPa → **0.01 hPa**



**Wave-like bias structure
now over the Arctic**



Tropical Tropopause



Systematic Difference in **Tropical Tropopause** Region
essentially **disappeared** with **new ECMWF scheme**


Borsche et al. 2007, *Geophys. Res. Lett.*

More on the Tropical Tropopause at **9:50**




Observing Climate

Preliminary Results for FORMOSAT-3/COSMIC



Foelsche et al. 2007, *TAO*, submitted

Monthly Mean Dry Temperature




Monthly Mean Dry Temperature

FORMOSAT-3/COSMIC August 2006


All satellites

Aug2006: Systematic Dry Temp Difference ECMWF-COSMIC




Seasonal Mean Dry Temperature

SON 2006 COSMIC-FM1 minus CHAMP



COSMIC Sampling


Sep 2006 : COSMIC-C1 Event Distribution



Seasonal Mean Dry Temperature

SON 2006 COSMIC-FM1 minus CHAMP


SON 2006: (COSMIC-C1- Sampl Err) - (CHAMP-Sampl Err)



Consistency of RO Data


SON 2006 **COSMIC-FM5** minus **CHAMP**

SON 2006: (FM5- Sampl Err) - (CHAMP-Sampl Err)




Consistency of RO Data

SON 2006 COSMIC-FM1 **minus** COSMIC-FM3




Consistency of RO Data

SON 2006 COSMIC-FM1 minus COSMIC-FM3



System. Difference – July 2007

December 2006: **Assimilation of RO data at ECMWF**





System. Difference – July 2007




Wavelike bias **disappeared** with **assimilation of RO data**

Systematic differences **above 30 km** considerably reduced

Sampling Error – July 2007


Azimuth angle further opened

July 2007: COSMIC Dry Temperature Sampling Error




Consistency – July 2007

COSMIC-FM1 **minus** COSMIC-FM2



Consistency – July 2007

COSMIC-FM2 minus COSMIC-FM4





Conclusions



Radio occultation measurements have a great potential for **monitoring of the global climate**

Comparison with existing climatologies:

- Deficiencies in ECMWF (Ant)arctic **winter stratosphere** disappeared with assimilation of RO data
- CHAMP **tropical tropopause** consistently **warmer** than ECMWF but only until February 2006
- Differences above 30 km: data from future missions (**MetOp**)

Comparison CHAMP – FORMOSAT-3/COSMIC:

- Preliminary but very promising: **consistency**

The FORMOSAT-3/COSMIC teams are especially acknowledged for the early provision of data!



Outlook



Initialization of RO profiles after Dec 2006 with ECMWF **forecasts** instead of **analyses** (since the analyses contain RO data now)

Measure of the degree of influence of the **background** provided

Cross Validation of RO climatologies from different satellites (especially including **MetOp**)



Challenges



- High altitude initialization
- Residual Ionospheric errors
- Potential reasons for common systematic errors
- Independent validation data
- SI Traceability
- Continuation of (dedicated) RO missions



Thank you!